

**POST-OPERATIVE REHABILITATION PROTOCOL  
ARTHROSCOPIC SUBACROMIAL DECOMPRESSION, DISTAL CLAVICLE  
EXCISION**



TODD C. MOEN, M.D.  
9301 N. Central Expressway, Suite 400  
Dallas, Texas 75231  
(214) 220-2468

Contact: Jennifer Kedroski, Dr. Moen's Coordinator, extension 3352

**RATIONALE:**

This protocol is designed with the goals of: 1) restoring pain-free motion, 2) restoring pain-free function. This protocol emphasizes a self-directed home exercise program. The Therapist and patient will determine the optimum interval and frequency of visits based on individual patient needs.

**IT IS CRITICALLY IMPORTANT THAT THE PATIENT PERFORM THE EXERCISES AS DIRECTED. NOT PERFORMING THE EXERCISES WILL LEAD TO A SUB-OPTIMAL RESULT FROM THE SURGERY.**

Exercises are best performed 2 - 3 times per day, for a 5 – 10 minutes at a time; they are more effective when performed in short bursts rather than a single longer session.

Weeks 1 - 6

**PRECAUTIONS:**

- No strengthening exercises
- No lifting objects > 5 lbs
- All range of motion exercises should be progressed as tolerated, as post-operative swelling and discomfort allow
- Emphasize slow, steady improvement each and every day – do not attempt large “heroic” improvement on any one given day.

**EXERCISES:**

- Pendulum/Codman's exercises for dressing activities, personal hygiene
- Passive Range of motion exercises, all planes
- Active Assist range of motion exercises, progress to Active ROM as tolerated; emphasize early use of pulley exercises to facilitate
- Scapular Retraction with arms at side and
- Scapular Kinematics exercises

- Elbow, wrist, finger exercises

Weeks 6 – 12

EXERCISES:

- Continue passive, active ROM exercises as necessary to achieve full ROM
- Progress from active assist to active ROM exercises as tolerated. Progress Active ROM as tolerated.
- Continue scapular retraction and kinematic exercises
- ***Emphasize appropriate scapular retraction and kinematics as active motion progresses***
- Begin Progressive Strengthening Program
- Begin with theraband exercises.
- Progress from therabands to light weights, and then progress as tolerated
- Goal of Progressive Strengthening program is to be performing strengthening exercises as tolerated in a gym by 4 ½ months post-op
- Theraband strengthening
- Low row exercises